arXiv:1209.6093v1 [hep-ph] 26 Sep 2012

BU-HEPP-12-03

# The Thomas-Fermi Quark Model: Non-Relativistic Aspects

Quan Liu, Walter Wilcox

*Department of Physics, Baylor University, Waco, TX 76798-7316*

Non-relativistic aspects of the Thomas-Fermi statistical quark model are developed. A review is given and our modified approach to spin in the model is explained. Our results are limited so far to two inequivalent simultaneous wave functions which can apply to multiple degenerate flavors. An explicit spin interaction is introduced, which requires the introduction of a generalized spin "flavor". Although the model is designed to be most reliable for many-quark states, we find surprisingly that it may be used to fit the low energy spectrum of octet and decouplet baryons. The low energy fit allows us to investigate the six-quark doubly strange $H$-dibaryon state, possible 6 quark nucleon-nucleon resonances and flavor symmetric strange states of higher quark content.

PACS numbers: 12.39.-x,12.39.Mk,12.40.Ee,21.65.Qr

## I. INTRODUCTION

The theoretical aim of our Thomas-Fermi ("TF") statistical quark model, introduced in Ref.[1], was to explore many-quark baryon states including strage matter[2, 3]. It is a useful new tool for the quark physicist to quickly assess the possibility of bound and resonance states in preparation for much more detailed and expensive lattice QCD calculations. The TF semi-classical model combines Coulombic quark interactions with a bag model type spherical confinement assumption. The model was seen to be extremely versatile in that a number of widely varying physics scenarios, including non-relativistic, extreme relativistic, massive gluon, or color-flavor locking, can be addressed. The development in Ref.[1], however, was limited to an investigation of a single quark wave function representing an equal number of $N_f$ mass-degenerate flavors. We would like to build upon our previous results to show how wave functions for unequal numbers of degenerate flavors or non-degenerate masses can be determined though the coupled differential equations. Although the model is designed to be most reliable for many-quark states, we find surprisingly that it may be used to fit the entire low energy spectrum of baryons. In order to do so, the model had to be extended to include an explicit spin splitting term. We introduce spin as a generalized "flavor" in our non-relativistic model. This change keeps the quark differential equation unmodified and just induces an overall scale change in the energy expressions. The non-relativistic nature of the model is a shortcoming, but we view our work as the appropriate initial step toward developing a more realistic relativistic version of the model. However, we note that the quark mass parameters dominate the overall baryon masses and that the overall fits are extremely good, so the model is self-consistent.

In the next section we will describe some of the changes in the formalism necessary to introduce quark spin as a generalized flavor. We will then present the differential equations and the generalized expressions for the quark energies in Section III for states with two different wave functions. More of the formalism for the consideration of spin will then be presented in Section IV. We will describe the computer program that solves the TF wave function differential equations and present our fit to the baryon spectrum in Section V. Note that this model is designed only to look for true many-quark bound states and resonances rather than bound nucleonic states like the deuteron. In this sense, the model complements calculations of loosely bound states as in Ref.[5]. As an application, we will attempt such a search in Section VI A for the hypothesized $H$-dibaryon state[4], make connection to nucleon-nucleon resonances in SectionVI B, and apply the formalism to high multi-quark strange states in SectionVI C.

## II. MATHEMATICAL PRELIMINARIES

Let us recall how the TF quark model is formulated. The original flavor and color number, $n_i^I(p_F)$, and non-interacting quark kinetic energy, $\mathcal{E}_i^I(p_F)$, densities were

$$n_i^I(p_F) = 2\int^{p_F} \frac{d^3p_i^I}{(2\pi\hbar)^3} = \frac{((p_F)_i^I)^3}{3\pi^2\hbar^3}, \tag{1}$$

and

$$\mathcal{E}_i^I(p_F) = 2\int^{p_F} \frac{d^3p_i^I}{(2\pi\hbar)^3}\frac{(p_i^I)^2}{2m} = \frac{(3\pi^2\hbar^3 n_i^I(p_F))^{5/3}}{10\pi^2\hbar^3 m}, \tag{2}$$

where $(p_F)_i^I$ is the Fermi momenta and where the "$I$" superscript stands for flavor and the "$i$" subscript stands for color. Notice the explicit degenerate spin factors of 2. Such an assumption of degeneracy is excellent for atomic physics, marginal for nuclear physics, but unrealistic for hadronic physics. The addition of a spin term in the TF Hamiltonian requires the introduction of non-equal spin densities. Instead, we will define new quantities distinguished by caligraphic superscripts,

$$n_i^{\mathcal{I}}(p_F) \equiv \int^{p_F} \frac{d^3p_i^{\mathcal{I}}}{(2\pi\hbar)^3} = \frac{((p_F)_i^{\mathcal{I}})^3}{6\pi^2\hbar^3}, \tag{3}$$

$$\mathcal{E}_i^{\mathcal{I}}(p_F) \equiv \int^{p_F} \frac{d^3p_i^{\mathcal{I}}}{(2\pi\hbar)^3}\frac{(p_i^{\mathcal{I}})^2}{2m} = \frac{(6\pi^2\hbar^3 n_i^{\mathcal{I}}(p_F))^{5/3}}{20\pi^2\hbar^3 m}, \tag{4}$$

where the spin specification is now included in an augmented flavor-spin index, $\mathcal{I}$. After variation of the densities in the energy functional, Eq.(21) of Ref.[1] still holds:

$$\frac{(p_F^{\mathcal{I}})^2}{2m^I} = -\lambda^{\mathcal{I}} + \frac{3\times\frac{4}{3}g^2}{(3A-1)}\left(\frac{N_{\mathcal{I}}-1}{N_{\mathcal{I}}}\int^{r_{max}} d^3r'\,\frac{N_{\mathcal{I}}(r')}{|\vec{r}-\vec{r}\,'|} + \sum_{J\neq I}\int^{r_{max}} d^3r'\,\frac{n^{\mathcal{J}}(r')}{|\vec{r}-\vec{r}\,'|}\right).$$

However, the previous relationship between the TF spatial wave function and the density,

$$f^I(r) \equiv \frac{ra}{2\times\frac{4}{3}\alpha_s}(3\pi^2 n^I(r))^{2/3}. \tag{5}$$

is now

$$f^{\mathcal{I}}(r) \equiv \frac{ra}{2\times\frac{4}{3}\alpha_s}(6\pi^2 N_{\mathcal{I}}(r))^{2/3}. \tag{6}$$

Rather than change the TF differential equations it turns out that all one need do to accommodate the new approach is to make a change in the overall spatial scale factor. The previous definition

$$r = Rx, \qquad R \equiv \left(\frac{a}{2\times\frac{4}{3}\alpha_s}\right)\left[\frac{3\pi A}{4}\right]^{2/3}, \tag{7}$$

is now replaced by

$$r = \mathcal{R}x, \qquad \mathcal{R} \equiv \left(\frac{a}{2 \times \frac{4}{3}\alpha_s}\right)\left[\frac{3\pi A}{2}\right]^{2/3}. \tag{8}$$

equivalent to a simple change in the underlying scale factor:

$$a \longrightarrow 2^{2/3}a. \tag{9}$$

The form of the differential TF equations in Ref.[1] and the wave function normalization,

$$\int_0^{x_{max}} dx\sqrt{x}(f^{\mathcal{I}}(x))^{3/2} = \frac{N_{\mathcal{I}}}{3A}. \tag{10}$$

are unchanged from before.

## III. GENERALIZED MODEL FOR TWO WAVE FUNCTIONS

In Ref.[1], a non-relativistic model with equal numbers of $N_f$ degenerate flavors was considered in a case study. However, in order to do realistic spectrum calculations, we have to generalize to unequal numbers of flavors. In this section, we explain how to solve for systems with 2 separate masses or systems with two equal masses, but different flavor amounts (numbers). We will see this is equivalent to the introduction of two quark flavor-degeneracy factors, $(N_f)_1 \equiv g_1$ and $(N_f)_2 \equiv g_2$. The differential form of the TF quark equations for this case, which involves two distinct wave functions, is

$$\alpha_{\mathcal{I}}\frac{d^2 f^{\mathcal{I}}(x)}{dx^2} = -\frac{A}{A-\frac{1}{3}}\frac{1}{\sqrt{x}}\left(\frac{N_{\mathcal{I}}-1}{N_{\mathcal{I}}}\left(f^{\mathcal{I}}(x)\right)^{\frac{3}{2}} + \sum_{\mathcal{J}\neq\mathcal{I}}\left(f^{\mathcal{I}}(x)\right)^{\frac{3}{2}}\right). \tag{11}$$

Here $\alpha_{\mathcal{I}}$ is the mass ratio of the lightest mass quark, $m_1$, to quark flavor $\mathcal{I}$,

$$\alpha_{\mathcal{I}} = \frac{m_1}{m_{\mathcal{I}}}. \tag{12}$$

and $N_{\mathcal{I}}$ is the number of quarks with generalized flavor $\mathcal{I}$. (Note we are using subscripts on $N_{\mathcal{I}}$, $\alpha_{\mathcal{I}}$ and $m_{\mathcal{I}}$ so that they will not be mistaken for powers. This is in contrast to Ref.[1] all these quantities were denoted with superscripts.) For $g_1$ flavors with equal number $N_1$, considered the state with the larger radius, $x_1 \geq x_2$, and $g_2$ flavors with equal numbers $N_2$, we have

$$g_1N_1 + g_2N_2 = 3A, \tag{13}$$

for quark quantum numbers, and the generalized two wave function equations are:

$$\alpha_1\frac{d^2 f^1(x)}{dx^2} = -\frac{A}{A-\frac{1}{3}}\frac{1}{\sqrt{x}}\left[\left(\frac{N_1-1}{N_1}+g_1-1\right)\left(f^1(x)\right)^{3/2} + g_2\left(f^2(x)\right)^{3/2}\right], \tag{14}$$

$$\alpha_2\frac{d^2 f^2(x)}{dx^2} = -\frac{A}{A-\frac{1}{3}}\frac{1}{\sqrt{x}}\left[\left(\frac{N_2-1}{N_2}+g_2-1\right)\left(f^2(x)\right)^{3/2} + g_1\left(f^1(x)\right)^{3/2}\right]. \tag{15}$$

We assume there is a linear relation between $f^1(x)$ and $f^2(x)$: $f^1(x) = kf^2(x)$. The equations (14) and (15) become:

$$\alpha_1 k \frac{d^2 f^2(x)}{dx^2} = -\frac{A}{A-\frac{1}{3}} \frac{1}{\sqrt{x}} \left[ \left( \frac{N_1 - 1}{N_1} + g_1 - 1 \right) k^{3/2} + g_2 \right] \left(f^2(x)\right)^{3/2}, \tag{16}$$

$$\alpha_2 \frac{d^2 f^2(x)}{dx^2} = -\frac{A}{A-\frac{1}{3}} \frac{1}{\sqrt{x}} \left[ \left( \frac{N_2 - 1}{N_2} + g_2 - 1 \right) + g_1 k^{3/2} \right] \left(f^2(x)\right)^{3/2}. \tag{17}$$

To make these two equation consistent with each other implies that

$$\Rightarrow \alpha_1 g_1 k^{5/2} - \alpha_2 \left( \frac{N_1 - 1}{N_1} + g_1 - 1 \right) k^{3/2} + \alpha_1 \left( \frac{N_2 - 1}{N_2} + g_2 - 1 \right) k - \alpha_2 g_2 = 0. \tag{18}$$

We solve this condition numerically for the $k$ value for each particle parameter set.

Remember we have two normalization conditions:

$$\begin{cases} \int_0^{x_2} \sqrt{x} \left(f^2(x)\right)^{3/2} dx = (N_2/3A), \\ \int_0^{x_1} \sqrt{x} \left(f^1(x)\right)^{3/2} dx = (N_1/3A). \end{cases} \tag{19}$$

With Equation (19), we can express our normalization conditions in the form of boundary conditions:

$$\begin{cases} \left( x\frac{df^2(x)}{dx} - f^2(x) \right) |_{x_2} = -\frac{A}{A-\frac{1}{3}} \left[ g_1 k^{3/2} + \left( \frac{N_2-1}{N_2} + g_2 - 1 \right) \right] \frac{N_2}{3A\alpha_2}, \\ \left( x\frac{df^1(x)}{dx} - f^1(x) \right) |_{x_1} = -\frac{1}{\alpha_1}. \end{cases} \tag{20}$$

The $f^1(x) = kf^2(x)$ condition must be consistent with the normalization conditions. Because of the attractive nature of the system, it is a natural assumption that the wave functions are compacted around the origin. That is, we assume no voids in the particle interiors. Although this is a natural assumption, we have not proven these configurations have the lowest energies.

With these modified boundary conditions, we can derive the expression for kinetic, $T$, and potential, $U$, energies:

$$T = \sum_I \frac{12}{5\pi} \left( \frac{3\pi A}{2} \right)^{1/3} \frac{\frac{4}{3}g^2 \times \frac{4}{3}\alpha_s}{a} \int_0^{x_I} \frac{\left(f^I(x)\right)^{5/2}}{\sqrt{x}} dx \tag{21}$$

$$= \frac{12}{5\pi} \left( \frac{3\pi A}{2} \right)^{1/3} \frac{\frac{4}{3}g^2 \times \frac{4}{3}\alpha_s}{a} \left[ g_1\alpha_1 \int_0^{x_1} \frac{\left(f^1(x)\right)^{5/2}}{\sqrt{x}} dx + g_2\alpha_2 \int_0^{x_2} \frac{\left(f^2(x)\right)^{5/2}}{\sqrt{x}} dx \right] \tag{22}$$

$$\begin{aligned} &= \frac{12}{5\pi} \left( \frac{3\pi A}{2} \right)^{1/3} \frac{\frac{4}{3}g^2 \times \frac{4}{3}\alpha_s}{a} \left[ -\frac{5}{7} g_2 \frac{df^2(x)}{dx}|_{x_2} \frac{N_2}{3A} \left( \alpha_2 - \frac{k g_1 \alpha_1}{\frac{N_1-1}{N_1} + g_1 - 1} \right) \right. \\ &\left. -\frac{5}{7} \frac{\frac{A-\frac{1}{3}}{A} g_1 \alpha_1}{\frac{N_1-1}{N_1} + g_1 - 1} \frac{df^1(x)}{dx}|_{x_1} + \frac{4}{7} g_2 \alpha_2 \left(f^2(x_2)\right)^{5/2} \sqrt{x_2} + \frac{4}{7} g_1 \alpha_1 \left(f^1(x_1)\right)^{5/2} \sqrt{x_1} \right]. \end{aligned} \tag{23}$$

$$
\begin{aligned}
U = &-\frac{2}{\pi}\frac{A}{A-\frac{1}{3}}\left(\frac{3\pi A}{2}\right)^{1/3}\frac{\frac{4}{3}g^2\times\frac{4}{3}\alpha_s}{a} \\
&\times\left[\sum_I \frac{N^I-1}{N^I}\left[\int_0^{x_{max_I}} dx\frac{\left(f^I(x)\right)^{4/3}}{\sqrt{x}}\int_0^x dx'\sqrt{x'}f^I(x') + \int_0^{x_{max_I}} dx\sqrt{x}f^I(x)\int_x^{x_{max_I}} dx'\frac{\left(f^I(x')\right)^{4/3}}{\sqrt{x'}}\right]\right. \\
&\left.+\sum_{I\neq J}\left[\int_0^{x_{max_I}} dx\frac{\left(f^I(x)\right)^{4/3}}{\sqrt{x}}\int_0^x dx'\sqrt{x'}f^J(x') + \int_0^{x_{max_I}} dx\sqrt{x}f^I(x)\int_x^{x_{max_J}} dx'\frac{\left(f^J(x')\right)^{4/3}}{\sqrt{x'}}\right]\right]
\end{aligned}
\tag{24}
$$

$$
\begin{aligned}
= &-\frac{2}{\pi}\left(\frac{3\pi A}{2}\right)^{1/3}\frac{\frac{4}{3}g^2\times\frac{4}{3}\alpha_s}{a}\times\left[\frac{N_2}{3A}\frac{df^2(x)}{dx}|_{x_2}\left(-\frac{12}{7}\alpha_2 g_2+\frac{12}{7}\frac{g_1g_2k\alpha_1}{\frac{N_1-1}{N_1}+g_1-1}\right)\right. \\
&+\frac{df^1(x)}{dx}|_{x_1}\left(-\frac{g_1g_2\alpha_1}{\frac{N_1-1}{N_1}+g_1-1}\frac{N_2}{3A}-k^{3/2}g_1\alpha_1\frac{N_2}{3A}-\frac{5}{7}\frac{\frac{A-1/3}{A}\alpha_1g_1}{\frac{N_1-1}{N_1}+g_1-1}\right) \\
&+\frac{\frac{A-1/3}{A}\alpha_1g_1}{\frac{N_1-1}{N_1}+g_1-1}-\frac{N_2}{3A}\alpha_1g_1k^{3/2}-\frac{g_1g_2\alpha_1}{\frac{N_1-1}{N_1}+g_1-1}\frac{N_2}{3A} \\
&\left.+\frac{4}{7}g_2\alpha_2\left(f^2(x_2)\right)^{5/2}\sqrt{x_2}+\frac{4}{7}g_1\alpha_1\left(f^1(x_1)\right)^{5/2}\sqrt{x_1}\right].
\end{aligned}
\tag{25}
$$

Of course, one is not limited to two different wave functions in this model, and the above equations can be generalized. Such a generalization is not necessary for the low energy baryon spectrum.

## IV. MORE ON SPIN

As outlined above, the traditional TF model simply assumes a spin degeneracy factor of 2 in Eqs.(1) and (2). We initially attempted to develop such a model based upon Ref.[1]. In this traditional approach there is an intrinsic splitting between states like the nucleon and $\Delta^{++}$ (or $\Delta^{-}$) already just from the different TF flavor wave functions, exclusive of spin, with the correct ordering of states. That is, there is a degeneracy splitting from the different quark flavor content, *uud* versus *uuu*, for example. However, the nucleon-delta splitting turns out to be much smaller than the actual splitting for reasonable parameter values. In our attempts to fit the low energy baryon spectrum, we found that our $\alpha_s$ and $m_1$ parameters (the latter sets the overall scale) were driven to unrealistic values to try to account for such splittings. The model was incapable of producing a realistic low energy spectrum. It is necessary to add an explicit spin interaction term.

There is a problem of course because products of the spin "up" and "down" (usually taken along the z-axis) in the semi-classical TF model do not combine to form "good" total angular momentum states. Non-relativistic baryon states such as the proton and neutron have total angular momentum $J=\frac{1}{2}$ and magnetic quantum number, $m_J$. To explain our approach to incorporating spin, consider the appropriately symmetrized proton flavor-spin $J=\frac{1}{2}$, $m_J=\frac{1}{2}$ wave function:

$$|P,+>\equiv|uud>(2|++->-|+-+>-|-++>)/(3\sqrt{2})+\text{cyclic permutations}. \qquad (26)$$

The TF quark model is incapable of reproducing this linear combination. Instead given this wave function the TF model simply asks the probabilities are of certain configurations. In the proton world, the possible spin up configurations are:

$$(u^\uparrow u^\uparrow d^\downarrow), \quad (u^\uparrow u^\downarrow d^\uparrow).$$

The proton is then in the configuration:

$$\frac{2}{3}(u^\uparrow u^\uparrow d^\downarrow)+\frac{1}{3}(u^\uparrow u^\downarrow d^\uparrow).$$

(We assign no meaning to the flavor or spin sequential ordering.) It is the configurations which have distinct masses in the TF model. Thus, the TF spin model deals with probabilities of configurations rather than spin amplitudes. We assume the mass of the physical state is the probability weighted average of the configurations. Using the flavor degeneracy factors introduced in the last section, these two configurations are classified:

$$(uud): \; g_1=1, N_1=2; \, g_2=1, N_2=1.$$
$$(uud): \; g_1=3, N_1=1; N_2=0.$$

That is, the first configuration has two identical particles (the $u$ quarks) and a second generalized flavor ($d$), whereas the second configuration simply has 3 non-identical but mass-degenerate quarks, with no second set of particle labelings necessary; note how the different spin $u$ quarks are treated as different flavors. We will have more to say on the violation of rotational symmetry later in this section.

The classical interaction of a magnetically charged particle in an external magnetic field is given by

$$(H_m^{class})_{ij}=-\vec{m}_i\cdot\vec{B}_j, \qquad (27)$$

where $\vec{m}_i$ is the magnetic moment of particle $i$ and $\vec{B}_j$ is the external magnetic field of particle $j$. A magnetic dipole field is given by

$$\vec{B}^d=\frac{3\hat{r}(\hat{r}\cdot\vec{m})-\vec{m}}{r^3}+\frac{8\pi}{3}\vec{m}\,\delta(\vec{r}), \qquad (28)$$

where $\delta(\vec{r})$ is a Dirac delta function. In our case, the spherical symmetry of all integrations require the interaction to be of the form,

$$(H_m^{class})_{ij} = -\frac{8\pi}{3}\vec{m}_i \cdot \vec{m}_j \, \delta(\vec{r}). \tag{29}$$

These considerations lead us to postulate the form of the color magnetic interaction in our model:

$$(H_m)_{ij} = -\frac{8\pi}{3}\gamma_i\gamma_j(S_z^{\mathcal{I}})_i(S_z^{\mathcal{J}})_j(\vec{q}_i \cdot \vec{q}_j)\, \delta(\vec{r}), \tag{30}$$

where the superscripts $\mathcal{I}$ and $\mathcal{F}$ are flavor labels and $i, j$ are particle number labels which will take on values from 1 to $N_{\mathcal{I}}$, $N_{\mathcal{J}}$. We define

$$\gamma_i \equiv \frac{\mathrm{g}}{2m_i c}, \tag{31}$$

where g is the gyromagnetic factor, which takes on a value of 1 classically and a value of 2 to lowest order in QED perturbation theory for the electron and muon. Note that the specialization to spins along the z-axis is appropriate for our individual spin basis. For two quarks,

$$\vec{q}_i \cdot \vec{q}_j = \begin{cases} \frac{4}{3}g^2, \text{ same color} \\ -\frac{2}{3}g^2, \text{ different color} \end{cases} \tag{32}$$

The treatment of potential energy terms is explained in Ref.[1]. We sum over same flavor and different flavor contributions, using a single particle normalization for the particle densities ($\hat{n}^{\mathcal{I}}(r) = 3n^{\mathcal{I}}(r)/N_{\mathcal{I}}$). We then average over the color interactions in Eq.(30) using (32) and the color probabilities, $P_{ij}$. The result for the magnetic spin interaction energy, $E_m$, may be written:

$$\begin{aligned} E_m = \frac{8\pi}{3}\frac{9 \times \frac{4}{3}g^2}{(3A-1)} \Bigg[ \left(\frac{\hbar}{2}\right)^2 \sum_{\mathcal{I}} \gamma_{\mathcal{I}}^2 \left(\frac{N_{\mathcal{I}}-1}{2N_{\mathcal{I}}}\right) \int d^3r \, \left(n^{\mathcal{I}}(r)\right)^2 \\ + \sum_{\mathcal{I}<\mathcal{J}} \gamma_{\mathcal{I}}\gamma_{\mathcal{J}} \sum_{i=1}^{N_{\mathcal{I}}}\sum_{j=1}^{N_{\mathcal{J}}} (S_z^{\mathcal{I}})_i (S_z^{\mathcal{J}})_j \int d^3r \, \left(\frac{n^{\mathcal{I}}(r) n^{\mathcal{J}}(r)}{N_{\mathcal{I}}N_{\mathcal{J}}}\right) \Bigg]. \end{aligned} \tag{33}$$

Switching to the TF wave function, $f^{\mathcal{I}}(r)$ defined in Eq.6, and using the $x$ variable defined in Eq.8 now gives

$$\begin{aligned} E_m = \frac{16\hbar c}{3\pi^2(3A-1)} \frac{(\frac{4}{3}\alpha_s)^4}{a} \left(\frac{\mathrm{g}}{2}\right)^2 \Bigg[ \sum_{\mathcal{I}} \alpha_{\mathcal{I}}^2 \frac{N_{\mathcal{I}}-1}{2N_{\mathcal{I}}} \int dx \, \frac{(f^{\mathcal{I}}(x))^3}{x} \\ + \sum_{\mathcal{I}<\mathcal{J}} \alpha_{\mathcal{I}}\alpha_{\mathcal{J}} \sum_{i=1}^{N_{\mathcal{I}}}\sum_{j=1}^{N_{\mathcal{J}}} (\hat{S}_z^{\mathcal{I}})_i (\hat{S}_z^{\mathcal{J}})_j \int dx \frac{1}{x} \frac{(f^{\mathcal{I}}(x))^{3/2}}{N_{\mathcal{I}}} \frac{(f^{\mathcal{J}}(x))^{3/2}}{N_{\mathcal{J}}}. \Bigg] \end{aligned} \tag{34}$$

where the unit normalized normalized spins have $(\hat{S}_z^{\mathcal{J}})_j = 1$ for spin up and $(\hat{S}_z^{\mathcal{J}})_j = -1$ for spin down along z.

As pointed out above, the introduction of spin labeling as an extended flavor attribute has the obvious shortcoming of loss of rotational symmetry. One way of stating the issue is that there is a conflict between the configurations used to determine flavor content, which use an individual spin basis with good values of $J_z^i$, where $i$ labels the particles, and the rotationally invariant total spin states which have good total angular momentum quantum numbers. This problem will not affect multi-quark total spin 0 or 1/2 states. For example, the proton $m = \pm\frac{1}{2}$ states both project unambiguously to the configuration $\frac{1}{3}uud + \frac{2}{3}uud$. However, the spin $\frac{3}{2}$ $m = \frac{1}{2}$ state,

$$\frac{1}{\sqrt{3}}(| -++ > + | +-+ > + | ++- >),$$

produces a different mass than the $m = \frac{3}{2}$ state when projected into the individual spin basis states. The $m = \frac{1}{2}$ state corresponds to a $g_1 = 1, N_1 = 2; g_2 = 1, N_2 = 1$ configuration in our model, whereas the $m = \frac{3}{2}$ state corresponds to $g = 1, N = 3$. These have different masses.

We believe the best projection is always performed in a maximum $m$ states, $|J, m = J >$. The reason has to do with what we will call the "maximum compatibility" of the two sets of basis states. As an example, let us consider the spin $|\frac{3}{2}, \frac{3}{2} >$ 3 quark states. Here there is only one way to build the state, namely $|\frac{1}{2} >_1 |\frac{1}{2} >_2 |\frac{1}{2} >_3$ in the product spin basis. This state is also an eigenstate of total $\vec{J}^2$ and $J_z$. The expectation value of the commutator of $\vec{J}^2$ with the individual $J_z$ vanishes:

$$< \frac{3}{2}, \frac{3}{2} | [\vec{J}^2, J_z^{1,2,3}] | \frac{3}{2}, \frac{3}{2} >= 0. \tag{35}$$

This commutator does not vanish in the $J = \frac{3}{2}, m = \frac{1}{2}$ state. As a more general example, consider the $J = 0, 1, 2, 3$ states produced by coupling 6 quarks. The $J = 0$ states will have a rotationally invariant projection, whereas the $J = 3, m = 3$ state will be maximally compatible with the states with good individual $J_z^{1,\ldots,6}$ since the commutator will again vanish. The maximal $m$ states, $J = 2, m = 2$ and $J = 1, m = 1$ will also be the most compatible states to use with $J = 2$ and $J = 1$, although here the commutators will no longer vanish.

## V. GROUND STATE BARYON FITS

In order for the model we are presenting to be predictive, we need to fix the phenomenological parameters. These 5 parameters are:

$B$: "Bag" constant

$m_1$: light quark mass

$m_s$: strange quark mass

$\alpha_s$: strong coupling constant

g: color gyromagnetic factor

First, we must understand that the model itself is not well designed to fit low quark number states, just as the atomic version would be poorly constructed to fit the low atomic binding energy states. In fact, applying this model to a "gas" of three quarks would seem to be impossible. However, just like the atomic model, an unreasonable aspect of the mathematics of the TF model is that such a fit seems to be entirely reasonable. An aspect that helps is the fact that in the hadronic case there are both the octet and decuplet ground states energies available which may be used to help make the fit more robust; that is, there are more states available than parameters which need to be fit.

There are 13 types of base configurations and associated TF wave functions that are necessary to fit all the octet and decuplet states. Table 1 lists the wave functions present in the ground state hadrons as well as the particles which partly or wholly share that wave function. For four of these wave functions there are both light quark ($u$ and $d$) and strange quark combinations present.

In all cases listed in Table 1 except the ones with either $g_1 = 3$, $N_1 = 3$ or $N_s = 3$, the consistency condition, Eq.(18), does not formally have a solution when the $g$ value or $N$ value is equal to one. These cases are indicated in the Table with the designation $1^+$ to indicate that the solution to (18) is defined by an infinitesimal approach to that parameter from above. For example, in the case of the $f_2$ wave functions, which contribute to the energies of the $P$, $N$ and $\Delta^+$ states, we have the robust numerical condition,

$$f_2^{2l}(x) = 1.23162 f_2^l(x), \tag{36}$$

for the relationship between $f^1$ ($f_2^{2l}$) and $f^2$ ($f_2^l$) for $0 < x < x_2$ in the limit $N_2 \longrightarrow 1$ from above. We use this limit to *define* the value of the model for $N_2 = 1$ exactly. (Note that $f^{2l}(x) = 0$ for $x_2 < x < x_1$.)

Table 2 gives both the base particle configuration, to be used in conjunction with Table 1, as well as the actual formulas for the spin splittings from Eq.(34) above. Besides the proton and neutron, note that a number of particles share the same base particle energies before magnetic spin splitting energies are added, such as the $\Delta^{++,-}$, the $\Delta^{+,0}$, the $\Sigma^{+,-}$, the $\Sigma^{*+,-}$, the $\Xi^{0,-}$, the

$\Xi^{*0,-}$, as well as the $\Lambda$, $\Sigma^0$ and $\Sigma^{*0}$. We list only the $\Delta^{++}$, $\Delta^+$, $\Sigma^+$, $\Sigma^{*+}$, $\Xi^0$, and $\Xi^{*0}$ particles in the Table because of these degeneracies, although the $\Lambda$, $\Sigma^0$ and $\Sigma^{*0}$ all get different masses after spin interaction.

There is a subtlety concerning the $\Xi^{0,-}$ and $\Xi^{*0,-}$ particles which prevents us from including them in our numerical results. All these particles contain two strange quarks and a light quark. In our numerical simulations we find that the two strange quark wave function,$f_6^s$ or $f_7^s$, has a smaller radius than the single light quark, $f_6^l$ or $f_7^l$, which in Table 1 is characterized as either $g_1 = 1^+, N_1 = 1$ or $g_1 = 1, N_1 = 1^+$. This means that the right hand side of Eq.(14) is zero for the region $x_2 < x < x_1$, where $f^2(x) = 0$, and that the light quark is "free" with no restoring force. The equations can still be formally solved and the normalization condition for $f^1$ in Eq.(19) can still be fulfilled, but we reject this solution as unrealistic. Thus, although the $\Xi^{0,-}$ and $\Xi^{*0,-}$ wave functions are formally calculable, we don't believe the model can not give correct wave functions in this case, and they are omitted from our numerical results.

A computer program in Mathematica has been developed and the parameters which fit the low energy baryon spectrum have been determined. Using a grid search algorithm we have found a preliminary best fit with the values: $B^{1/4}$=80 MeV, $m_1$=300 MeV/c$^2$, $m_s = 520$ MeV, $\alpha_s = 0.44$, g = 7.31. These values are phenomenologically appropriate and reasonable. The bag constant is slightly lower than in other studies, which means the non-relativistic fit is actually more self-consistent. The strong coupling constant is consistent with a bag model designed for heavy-light systems[6]. The gyromagnetic factor, g, is large compared to the electrodynamic case ($\sim 2$), but not unreasonably so. We do a global fit over the 9 particles listed below. Our results so far for our mass fit in MeV/c$^2$ units is (experimental numbers in parentheses):

**Spin$\frac{1}{2}$:**

P,N: 966 (939)
$\Lambda$: 1135 (1116)
$\Sigma^{+,-}$: 1214 (1189,1197)
$\Sigma^0$: 1169 (1192)

**Spin $\frac{3}{2}$:**

$\Delta^{++,-}$: 1221 (1232)

$\Delta^{+,0}$: 1205 (1232)

$\Sigma^{*+,-}$: 1416 (1385)

$\Sigma^{*0}$: 1391 (1385)

$\Omega^{-}$: 1702 (1672)

One can see that the model overestimates the strong isospin breaking effects from the particle wave functions for the differently charged $\Delta$, $\Sigma$ and $\Sigma^*$ particles. We note that the $\Delta^{++,-}$ and $\Delta^{+,0}$ particles actually come from different base configurations; similarly for the $\Sigma^{+,-}$ and $\Sigma^0$ as well as the $\Sigma^{*+,-}$ and $\Sigma^{*0}$. There should also be such isospin breaking effects in nature from the exclusion principle, but clearly they are not as large as seen here. Although the fit is not as good as for a typical non-relativistic quark model, we would not expect it to be. As explained above, the known ground state baryons badly violate the many quark semi-classical assumption. Nevertheless, we are encouraged by the overall reasonableness of the fit.

Table 3 gives a full accounting of the various particle energies in our model. Notice that the final particle energies involved are consistent with the non-relativistic assumption partly there are large cancellations between the kinetic and potential energies. Also notice that the radii are relatively large compared, for example, to measured charged particle electromagnetic radii. This is a result of the fit with a smaller value of $B^{1/4}$ than most standard phenomenologies. This is also partly responsible for the largely non-relativistic nature of the fits.

We now turn to try applying the phenomenology developed here to quark configurations in a number of sectors.

| Base configuration | Particle(s) | Wave function | $g_1$ | $N_1$ | $g_2$ | $N_2$ | $g_s$ | $N_s$ |
|---|---|---|---|---|---|---|---|---|
| $u^\uparrow u^\downarrow d^\uparrow, d^\uparrow d^\downarrow u^\uparrow$ | P,N | $f_1$ | 3 | 1 | -- | -- | -- | -- |
| $u^\uparrow u^\uparrow d^\downarrow, u^\uparrow u^\uparrow d^\uparrow$<br>$d^\uparrow d^\uparrow u^\downarrow, d^\uparrow d^\uparrow u^\uparrow$ | P,N<br>$\Delta^+$ | $f_2^{2l}, f_2^l$ | 1 | 2 | 1 | $1^+$ | -- | -- |
| $u^\uparrow u^\uparrow u^\uparrow, d^\uparrow d^\uparrow d^\uparrow$ | $\Delta^{++}$ | $f_3$ | 1 | 3 | -- | -- | -- | -- |
| $u^\uparrow u^\downarrow s^\uparrow, d^\uparrow d^\downarrow s^\uparrow$<br>$u^\uparrow d^\uparrow s^\downarrow, u^\uparrow d^\downarrow s^\uparrow$<br>$u^\downarrow d^\uparrow s^\uparrow, u^\uparrow d^\uparrow s^\uparrow$ | $\Sigma^+$<br>$\Sigma^0$<br>$\Sigma^{*0}$<br>$\Lambda$ | $f_4^l, f_4^s$ | 2 | 1 | -- | -- | $1^+$ | 1 |
| $u^\uparrow u^\uparrow s^\downarrow, u^\uparrow u^\uparrow s^\uparrow$<br>$d^\uparrow d^\uparrow s^\downarrow, d^\uparrow d^\uparrow s^\uparrow$ | $\Sigma^+$<br>$\Sigma^{*+}$ | $f_5^l, f_5^s$ | 1 | 2 | -- | -- | 1 | $1^+$ |
| $s^\uparrow s^\downarrow d^\uparrow, s^\uparrow s^\downarrow d^\uparrow$ | $\Xi^0$ | $f_6^l, f_6^s$ | $1^+$ | 1 | -- | -- | 2 | 1 |
| $s^\uparrow s^\uparrow u^\downarrow, s^\uparrow s^\uparrow u^\uparrow$<br>$s^\uparrow s^\uparrow d^\downarrow, s^\uparrow s^\uparrow d^\uparrow$ | $\Xi^0$<br>$\Xi^{*0}$ | $f_7^l, f_7^s$ | 1 | $1^+$ | -- | -- | 1 | 2 |
| $s^\uparrow s^\uparrow s^\uparrow$ | $\Omega^-$ | $f_8$ | -- | -- | -- | -- | 1 | 3 |

Table 1: Configuration wave functions.

| **Base particle configeration** | **Magnetic Energy splitting** $C \equiv \frac{16\hbar c}{3\pi^2(3A-1)} \frac{(4/3\alpha_s)^4}{a} \left(\frac{g}{2}\right)^2$ |
|---|---|
| $P = 1/3(u^\uparrow u^\downarrow d^\uparrow) + 2/3(u^\uparrow u^\uparrow d^\downarrow)$ | $E_m^P = C\left\{1/3\left(-\int dx \frac{(f_1)^3}{x}\right) + 2/3\left(1/4\int dx \frac{(f_2^{2l})^2}{x} - \int dx \frac{(f_2^{2l} f_2^l)^{3/2}}{x}\right)\right\}$ |
| $\Lambda = u^\uparrow d^\downarrow s^\uparrow$ | $E_m^\Lambda = C\left\{-\int dx \frac{(f_4^l)^3}{x}\right\}$ |
| $\Sigma^+ = 1/3(u^\uparrow u^\downarrow s^\uparrow) + 2/3(u^\uparrow u^\uparrow s^\downarrow)$ | $E_m^{\Sigma^+} = C\left\{1/3\left(-\int dx \frac{(f_4^l)^3}{x}\right) + 2/3\left(1/4\int dx \frac{(f_5^l)^3}{x} - \alpha\int dx \frac{(f_5^l f_5^s)^{3/2}}{x}\right)\right\}$ |
| $\Sigma^0 = 1/3(u^\uparrow d^\downarrow s^\uparrow) + 2/3(u^\uparrow d^\uparrow s^\downarrow)$ | $E_m^{\Sigma^0} = C\left\{1/3\left(-\int dx \frac{(f_4^l)^3}{x}\right) + 2/3\left(\int dx \frac{(f_4^l)^3}{x} - 2\alpha\int dx \frac{(f_4^l f_4^s)^{3/2}}{x}\right)\right\}$ |
| $\Xi^0 = 1/3(s^\uparrow s^\downarrow u^\uparrow) + 2/3(s^\uparrow s^\uparrow u^\downarrow)$ | $E_m^{\Xi^0} = C\left\{1/3\left(-\alpha^2\int dx \frac{(f_6^s)^3}{x}\right) + 2/3\left(\alpha^2/4\int dx \frac{(f_7^s)^3}{x} - \alpha\int dx \frac{(f_7^l f_7^s)^{3/2}}{x}\right)\right\}$ |
| $\Delta^{++} = (u^\uparrow u^\uparrow u^\uparrow)$ | $E_m^{\Delta^{++}} = C\left\{1/3\left(\int dx \frac{(f_3)^3}{x}\right)\right\}$ |
| $\Delta^+ = (u^\uparrow u^\uparrow d^\uparrow)$ | $E_m^{\Delta^+} = C\left\{1/4\left(\int dx \frac{(f_2^{2l})^3}{x}\right) + \left(\int dx \frac{(f_2^{2l} f_2^l)^{3/2}}{x}\right)\right\}$ |
| $\Sigma^{*+} = u^\uparrow u^\uparrow s^\uparrow$ | $E_m^{\Sigma^{*+}} = C\left\{1/4\left(\int dx \frac{(f_5^l)^3}{x}\right) + \alpha\left(\int dx \frac{(f_5^l f_5^s)^{3/2}}{x}\right)\right\}$ |
| $\Sigma^{*0} = (u^\uparrow d^\uparrow s^\uparrow)$ | $E_m^{\Sigma^{*0}} = C\left\{\left(\int dx \frac{(f_4^l)^3}{x}\right) + 2\alpha\left(\int dx \frac{(f_4^l f_4^s)^{3/2}}{x}\right)\right\}$ |
| $\Xi^{*0} = (u^\uparrow s^\uparrow s^\uparrow)$ | $E_m^{\Xi^{*0}} = C\left\{\alpha^2/4\left(\int dx \frac{(f_7^s)^3}{x}\right) + \alpha\left(\int dx \frac{(f_7^l f_7^s)^{3/2}}{x}\right)\right\}$ |
| $\Omega^- = (s^\uparrow s^\uparrow s^\uparrow)$ | $E_m^{\Omega^-} = C\left\{1/3\left(\int dx \frac{(f_8)^3}{x}\right)\right\}$ |

Table 2: Spin energy contributions.

| Particle | Radius(f) | Rest mass(MeV) | T(MeV) | U(MeV) | Spin(MeV) | Volume Energy(MeV) | Total(MeV) | Exp(MeV) |
|---|---|---|---|---|---|---|---|---|
| P | 1.52 | 900 | 201.34 | -141.46 | -67.21 | 73.67 | 966.34 | 938 |
| $\Delta^+$ | 1.52 | 900 | 218.08 | -134.74 | 143.01 | 78.73 | 1205.09 | 1232 |
| $\Delta^{++}$ | 1.72 | 900 | 231.72 | -124.18 | 99.50 | 114.26 | 1221.31 | 1232 |
| $\Lambda$ | 1.33 | 1121.74 | 185.36 | -147.26 | -77.16 | 52.04 | 1134.71 | 1115 |
| $\Sigma^+$ | 1.44 | 1121.74 | 216.90 | -145.18 | -40.49 | 61.41 | 1214.37 | 1194 |
| $\Sigma^0$ | 1.33 | 1121.74 | 185.36 | -147.26 | -42.80 | 52.04 | 1169.07 | 1192 |
| $\Sigma^{*+}$ | 1.44 | 1121.74 | 232.67 | -144.13 | 139.96 | 66.09 | 1416.31 | 1385 |
| $\Sigma^{*0}$ | 1.33 | 1121.74 | 185.36 | -147.26 | 179.94 | 52.04 | 1391.81 | 1385 |
| $\Omega^-$ | 1.51 | 1565.22 | 175.71 | -144.32 | 28.75 | 76.87 | 1702.22 | 1672 |

Table 3: The full set of octet and decuplet particle energies in the TF quark model.

## VI. THREE CASE STUDIES

In this section we will apply the fit found in the previous section to begin to explore some of the interesting phenomenology of high number multi-quark states. In particular, we will examine three possibilities: the $H$-dibaryon, nucleon-nucleon 6 quark resonances and high multi-quark strange states.

### A. $H$-Dibaryon Considerations

The $H$-dibaryon, which is an isospin $I = 0$, total angular momentum $J = 0$ $uuddss$ flavor state, is an interesting application of the model. Although there are many lattice results[7–14], there are systematic problems in the calculations due to lattice volume and quark mass effects. There are experimental results[15, 16] which limit the bound state energy to shallow values if indeed it exists at all[17]. The lattice results must be extrapolated in quark mass, and the results of this procedure are still rather uncertain[7]. Our model is designed to evaluate states for further lattice study, so it is interesting to study this possible bound state with the new tool we now possess, even though we are doing this study "post-lattice" at this point.

We may form a total $I = 0$, $J = 0$ combination from combining two particles in the $(\Lambda\Lambda)$, $(\Sigma\Sigma)$ or $(\Xi N)$ systems[5]. The individual particle isospins are $I_\Lambda = 0$, $I_\Sigma = 1$, $I_N = \frac{1}{2}$ and $I_\Xi = \frac{1}{2}$. One can look up the Clebsch-Gordon coefficients to couple these systems to a $|I_{tot} = 0>$ state. In quantum mechanics the states mix with one another, giving a coupled channel problem[8]. Such considerations are beyond the present model's purview. However, given the quantum wave functions each of these combinations may be examined separately for the lowest mass.

The possible TF quark 4 quark $uuddss$ $J = 0$ combinations are only three in number (we don't list some redundant configurations in the $m_u = m_d$ isospin limit):

$$(u^\uparrow u^\uparrow d^\downarrow d^\downarrow s^\uparrow s^\downarrow) : N_1 = 2, g_1 = 2, = 2; N_s = 1, g_s = 2,$$

$$(u^\uparrow u^\uparrow d^\uparrow d^\downarrow s^\uparrow s^\uparrow) : N_1 = 2, g_1 = 1; N_2 = 1, g_2 = 2; N_s = 2, g_s = 1,$$

$$(u^\uparrow u^\downarrow d^\uparrow d^\downarrow s^\uparrow s^\downarrow) : N_1 = 1, g_1 = 4; N_s = 1, g_s = 2.$$

Unfortunately, the $u, d$ asymmetric second combination, $(u^\uparrow u^\uparrow d^\uparrow d^\downarrow s^\uparrow s^\uparrow)$, can not be formed here because it involves three separate and different quark wave functions. This limits us to investigation of the $\Lambda\Lambda$ state only, although this is also likely the lightest state from Pauli blocking considerations.

Our result for the $\Lambda\Lambda$ $I = 0, J = 0$ state is 2360 MeV/c$^2$. The component energies and radius are:

$$\text{Quark rest mass} = 2244\,\text{MeV/c}^2,$$

$$\text{Kinetic energy} = 356\,\text{MeV/c}^2,$$

$$\text{Potential energy} = -266\,\text{MeV/c}^2,$$

$$\text{Volume energy} = 85.3\,\text{MeV/c}^2,$$

$$\text{Spin energy} = -58.5\,\text{MeV/c}^2,$$

$$\text{Radius} = 1.62\,\text{fm}$$

This is 90 MeV/c$^2$ more than the $\Lambda\Lambda$ threshold. Our present results indicate the $H$-dibaryon state is not bound in the TF quark model.

### B. Nucleon-Nucleon Resonances

When bound states are not present in a channel, our model is useful in predicting multi-quark baryon resonances. In Fig. 1 we show a "scan" of the configuration energies for the two wave function case for nucleon-nucleon scattering. Note these are the configuration energies, not the particle energies. To obtain particle resonance energies, we must proceed by projecting the quantum state into the particle configurations. Linear combinations of these configuration energies will be the actual resonance energies.

It turns out there are 8 different configuration in this case. Moving from left to right in the figure displays how the energy rises as more particle are put into a given flavor state. Given that $g_1N_1 + g_2N_2 = 6$, the explicit cases are:

$$N_1 = 0, ; N_2 = 2, g_2 = 3; \text{“}N_1.N_2\text{”} = 0.2$$

$$N_1 = 0, ; N_2 = 3, g_2 = 2; \text{“}N_1.N_2\text{”} = 0.3$$

$$N_1 = 0, ; N_2 = 6, g_2 = 1; \text{“}N_1.N_2\text{”} = 0.6$$

$$N_1 = 2, g_1 = 2; N_2 = 1, g_2 = 2; \text{“}N_1.N_2\text{”} = 2.1$$

$$N_1 = 3, g_1 = 1; N_2 = 1, g_2 = 3; \text{“}N_1.N_2\text{”} = 3.1$$

$$N_1 = 4, g_1 = 1; N_2 = 1, g_2 = 2; \text{“}N_1.N_2\text{”} = 4.1$$

$$N_1 = 4, g_1 = 1; N_2 = 2, g_2 = 1; \text{“}N_1.N_2\text{”} = 4.2$$

$$N_1 = 5, g_1 = 1; N_2 = 1, g_2 = 1; \text{“}N_1.N_2\text{”} = 5.1$$

Since each configuration is uniquely specified by the $N_1$ and $N_2$ values, we use these to numbers to form a fractional number defined by $N_1.N_2$ to plot the results in Fig. 1. The integer number represents the number of $N_1$ particles and the decimal number represents the number of $N_2$ particles. For example, 0.6 on the $N_1.N_2$ axis means $N_1 = 0; N_2 = 6, g_2 = 1$. Note that we only calculate the bare mass here in this quick scan of configuration energies.

The lowest configuration energy is seen to be in the $N_1.N_2 = 0.2$ state at 2016 MeV/c$^2$. This is as we expect from the exclusion principle since we are spreading 6 quarks across the maximum number of light flavors. In addition, the highest mass is $N_1.N_2 = 0.6$ at 2294 MeV/c$^2$. This is also as expected. We see that all of the states are greater than the two nucleon threshold and that there are no elemental six quark nucleon-nucleon bound states in our model. There is a resonance above two nucleon threshold in both *pp* and *np* scattering, but is impossible to tell if the states contributing are mesons and baryons or true 6 quark states.



FIG. 1: Nucleon-nucleon six quark configuration resonance chart. See the text for the meaning of the "$N_1.N_2$" notation.

### C. High Multi-Quark Strange States

We have studied the possibility of a bound $H$-dibaryon in Section VI A. Lattice results have already begun to accumulate for this system. We can get ahead of the lattice results by beginning to investigate the possibility of bound states with many more quarks.

We have investigated the possibility of addition $I = 0$ states constructed from $\Lambda$, $\Lambda$ wave functions. Such states can be thought of as multi-combinations of $H$-dibaryons with $12, 18, 24\ldots$ quarks. Such states would be very difficult if not impossible for the lattice to simulate. Here, it is a simple matter of changing the initial simulation parameters to accommodate these new states. In this model, we formally have a maximum of 4 light quark flavors. We have investigated the 12 quark configuration,

$$N_1 = 2, g_1 = 4; N_s = 2, g_s = 2,$$

the 18 quark configuration,

$$N_1 = 3, g_1 = 4; N_s = 3, g_s = 2,$$

and the 24 quark configuration,

$$N_1 = 4, g_1 = 4; N_s = 4, g_s = 2.$$

Higher quark number states are obvious generalizations. We find the following energies and masses in our investigation:

For the 12 quark case, we find the mass as 4777 MeV/c$^2$, 237 MeV/c$^2$, more than the 4 $\Lambda$ threshold and 57 MeV/c$^2$ more than twice the TF *H*-dibaryon mass. The component energies and radius are:

$$\begin{aligned}
&\text{Quark rest mass} \;=\; 4487\,\mathrm{MeV/c^2},\\
&\text{Kinetic energy} \;=\; 596\,\mathrm{MeV/c^2},\\
&\text{Potential energy} \;=\; -443\,\mathrm{MeV/c^2},\\
&\text{Volume energy} \;=\; 159\,\mathrm{MeV/c^2},\\
&\text{Spin energy} \;=\; -22.0\,\mathrm{MeV/c^2},\\
&\text{Radius} \;=\; 1.92\,\mathrm{fm}
\end{aligned}$$

For the 18 quark case, we find the mass as 7228 MeV/c$^2$, 417 MeV/c$^2$ more than the 6 $\Lambda$ threshold and 147 MeV/c$^2$ more than three times the TF *H*-dibaryon mass. The component energies and radius are:

$$\begin{aligned}
&\text{Quark rest mass} \;=\; 6731\,\mathrm{MeV/c^2},\\
&\text{Kinetic energy} \;=\; 861\,\mathrm{MeV/c^2},\\
&\text{Potential energy} \;=\; -601\,\mathrm{MeV/c^2},\\
&\text{Volume energy} = 251\,\mathrm{MeV/c^2},\\
&\text{Spin energy} = -13.1\,\mathrm{MeV/c^2},\\
&\text{Radius} \;=\; 2.24\,\mathrm{fm}
\end{aligned}$$

Finally, for the 24 quark case, we find the mass as 9679 MeV/c$^2$, 591 MeV/c$^2$ more than the 8 $\Lambda$ threshold and 239 MeV/c$^2$ more than four times the TF *H*-dibaryon mass. The component energies and radius are:

$$
\begin{aligned}
&\text{Quark rest mass} \;=\; 9874\,\mathrm{MeV/c^2},\\
&\text{Kinetic energy} \;=\; 1122\,\mathrm{MeV/c^2},\\
&\text{Potential energy} \;=\; -751\,\mathrm{MeV/c^2},\\
&\text{Volume energy} = \; 344\,\mathrm{MeV/c^2},\\
&\text{Spin energy} = \; -9.4\,\mathrm{MeV/c^2},\\
&\text{Radius} \;=\; 2.49\,\mathrm{fm}
\end{aligned}
$$

These states are moving further away from the bound particle thresholds. Interestingly, the systems are remaining relatively non-relativistic, with a smaller percentage of the component energy being kinetic as more quarks are added. In addition, the spin energies are becoming a smaller system component as well.

## VII. CONCLUSIONS

We have begun to develop the TF quark model as a tool in strong interaction dynamics. We have shown how to introduce spin into the TF quark model and found parameters which effectively fit the low energy baryon states. The results presented here show how to construct the necessary wave functions in the two wave function multi-flavor case with quantum numbers $N_1, g_1, N_2, g_2$.

We have undertaken three case studies: the *H*-dibaryon, nucleon-nucleon resonances and high multi-quark strange states. Our results for strange quark states show growing systems which are remaining relatively non-relativistic and whose spin interactions are a decreasing percentage of the total energy. There are no hints that the systems can be bound. However, these results should be regarded as initial indictions rather than final results. We have also confirmed a phenomenology in nucleon-nucleon scattering consistent with known low energy hadronic physics.

There are many remaining areas in which to continue the development. We note that the model may be extended:

- To be relativistic
- To include anti-quarks
- To include central heavy quarks
- To look at exotic forms of matter (eg., “strange” matter, color-flavor locked quarks from Cooper

pairs, or massive gluon models, or combinations)

The relativistic equation to be solved is given in Ref.[1], although the system energy formulas are not yet developed. We can not strongly defend our non-relativistic assumption other than pointing out that the present model should be regarded as a work in progress. The inclusion of anti-quarks would simply be accomplished by introducing a different color coupling for such particles and averaging over the couplings between quarks and anti-quarks. Heavy central quarks can be examined using Coulombic sources much like atomic TF systems. Perhaps most interestingly, the color-flavor locking scenario, partly examined in Ref.[1] can also be examined.

We believe that the TF quark model, in combination with lattice studies, can be an effective “tool” in evaluating characteristics of interesting baryonic states.

[1] W. Wilcox, Nucl. Phys. A **826** (1999) 49.

[2] E. Witten, Phys. Rev. D **30**: 272 (1984).

[3] E. Farhi, R. Jaffe, Phys. Rev. D **30**: 2379 (1984).

[4] R.L. Jaffe, Phys. Rev. **38** (1977) 195 [Erratum-ibid. **38** (1977) 617].

[5] J. Haidenbauer and U.-G. Mei$\beta$ner, Nucl. Phys. A **881** (2012) 44.

[6] W. Wilcox, Phys.Rev. D **31** (1985) 1081.

[7] W. Detmold *et al.*, AIP Conf. Proc. **1441**, pp. 332-334.

[8] T. Inuoe *et al.*, AIP Conf. Proc. **1441**, pp. 335-337.

[9] T. Sakai, K. Shimizu and K. Yazaki, Prog. Theor. Phys. Suppl. **137**, 121 (2000).

[10] I. Wetzorke and F. Karsch, Nucl. Phys. Proc. Suppl. **119**, 278 (2003).

[11] Z.H. Luo, M. Loan and X. Q. Luo, Mod. Phys. Lett. A22, 591 (2007).

[12] T. Inuoe *et al.*, Prog. Theor. Phys. **124**, 591 (2010).

[13] S.R. Beane *et al.*, Phys. Rev. Lett. **106**, 162001 (2011).

[14] T. Inuoe *et al.*, Phy. Rev. Lett. **106**, 162002 (2011).

[15] H. Takahashi *et al.*, Phys. Rev. Lett., **87**, 212502 (2001).

[16] C.J. Yoon *et al.*, Phy. Rev. C **75**, 02201 (2007).

[17] A. L. Trattner, PhD Thesis, LBL, UMI-32-54109 (2006).